Ikeda, K., Akiyama, H., Kondo, H. and Ikeda, K., Anti-tau-positive glial fibrillary tangles in the brain of postencephalitic parkinsonism of Economo type, 176
Imagawa, M., see Tanaka, H., 63
Inoki, R., see Uchida, Y., 114
Ishii, H., see Yoshida, M., 5

Jaffe, E.H., De Frias, V. and Ibarra, C., Changes in basal and stimulated release of endogenous serotonin from different nuclei of rats subjected to two models of depression, 157
Jellinger, K.A., see Bancher, C., 179
Johnson, J.A., see Tizzano, J.P., 12

Kamino, K., see Tanaka, H., 63
Katao, Y., see Minami, M., 161
Katsumata, S., see Minami, M., 161
Kawahara, H., Yoshida, M., Yokoo, H., Nishi, M. and Tanaka, M., Psychological stress increases serotonin release in the rat amygdala and prefrontal cortex assessed by in vivo microdialysis, 81
Keller, J.T., see Stanley, M.I., 89
Kent, T.A., see Stroemer, R.P., 51
Kiessling, M., see Gass, P., 39
Kim, H.J., see Chung, K., 85
Kimura, J., see Shimohama, S., 183
Kobayashi, H., see Tanaka, H., 63
Kobayashi, J.-i., Ohta, M. and Terada, Y., C fiber generates a slow Na^+ spike in the frog sciatic nerve, 93
Koda, H., see Mizumura, K., 75
Koga, C., see Yoshida, M., 5
Köhler, R., see Campos-Barros, A., 145
Kojima, T., see Tanaka, H., 63
Kondo, H., see Ikeda, K., 176
Krnjević, K., see Godfraind, J.M., 101
Kumazawa, T., see Mizumura, K., 75

Layrargues, G.P., see Mousseau, D.D., 192
Little, J., see Molchan, S.E., 187
Lovat, P.E., Pearson, A.D.J., Malcolm, A. and Redfern, C.P.F., Retinoic acid receptor expression during the in vitro differentiation of human neuroblastoma, 109
Lustig, H.S., see Ahern, K.v.B., 169

Maeda, S., see Uchida, Y., 114
Maekawa, K., see Minami, M., 161
Maeno, H., see Fujita, M., 59
Malcolm, A., see Lovat, P.E., 109
Manji, H., see Molchan, S.E., 187
Martínez, A., see de Rada, O.D., 121
Matsumoto, I., see Pujic, Z., 67
Matsumoto, S., see Shimohama, S., 183
Matsuya, T., see Uchida, Y., 114
May, T., 1-Methyl-4-phenylpyridinium (MPP^+) binds with high affinity to a β-carboline binding site located on monoamine oxidase type A in rat brain, 55
Mechoulam, R., see Nadler, V., 43
Meier, R.K., see Dieringer, N., 129
Meinhold, H., see Campos-Barros, A., 145
Miki, T., see Tanaka, H., 63
Minami, M., Hosoi, Y., Toya, T., Katao, Y., Maekawa, K., Katsumata, S., Yabuuchi, K., Onogi, T. and Satoh, M., In situ hybridization study of κ-opioid receptor mRNA in the rat brain, 161
Miyatake, T., see Tanaka, H., 63
Miyoshi, K., see Shimohama, S., 183
Mizoguchi, K., see Yoshida, M., 5
Mizumura, K., Koda, H. and Kumazawa, T., Augmenting effects of cyclic AMP on the heat response of canine testicular polymodal receptors, 75
Molchan, S.E., Manji, H., Chen, G., Dou, L., Little, J., Potter, W.Z. and Sunderland, T., Effects of chronic lithium treatment on platelet PKC isozymes in Alzheimer's and elderly control subjects, 187
Montuenga, L.M., see de Rada, O.D., 121
Morehead, M., see Sen, S., 117
Mousseau, D.D., Perney, P., Layrargues, G.P. and Butterworth, R.F., Selective loss of pallidal dopamine D_2 receptor density in hepatic encephalopathy, 192
Müller, F., see Campos-Barros, A., 145
Murphy, S., see Sen, S., 117
Murphy, S., see Bruner, G., 105

Na, H.S., see Chung, K., 85
Naash, M.I., see Peachey, N.S., 9
Nachemson, A.K. and Bennett, G.J., Does pain damage spinal cord neurons? Transsynaptic degeneration in rat following a surgical incision, 78
Nadler, V., Mechoulam, R. and Sokolovsky, M., The non-psychotropic cannabinoid (+)-(3*S*,4*S*)-7-hydroxy-Δ^6-tetrahydrocannabinol 1,1-dimethylheptyl (HU-211) attenuates *N*-methyl-D-aspartate receptor-mediated neurotoxicity in primary cultures of rat forebrain, 43
Nakano, I., see Tanaka, H., 63
Naruse, S., see Tanaka, H., 63
Nielander, H.B., French, P., Oestreicher, A.B., Gispen, W.H. and Schotman, P., Spontaneous morphological changes by overexpression of the growth-associated protein B-50/GAP-43 in a PC12 cell line, 46
Nishi, M., see Kawahara, H., 81
Nishimura, M., see Shimohama, S., 183
Nishimura, T., see Fujita, M., 59
Nishino, H., see Czurkó, A., 71
Nowak, J.Z. and Wawrocka, M., Intravitreal application of membrane-permeable analogs of cyclic GMP increases serotonin *N*-acetyltransferase (NAT) activity in retinas of light-exposed chicks: comparison with the effect of cyclic AMP analogs and darkness, 21
Nukina, N., see Tanaka, H., 63

Oestreicher, A.B., see Nielander, H.B., 46
Ohnishi, T., see Uchida, Y., 114
Ohta, M., see Kobayashi, J.-i., 93
Okamoto, K., Hirai, S., Amari, M., Watanabe, M. and Sakurai, A., Bunina bodies in amyotrophic lateral sclerosis immunostained with rabbit anti-cystatin C serum, 125
Onodera, O., see Tanaka, H., 63
Onogi, T., see Minami, M., 161

Pang, C.S., Brown, G.M., Tang, P.L. and Pang, S.F., G-protein linked melatonin binding sites in the chicken lung, 17
Pang, S.F., see Pang, C.S., 17
Panzica, F., see Franceschetti, S., 25
Park, M.J., see Chung, K., 85
Peachey, N.S., Goto, Y., Al-Ubaidi, M.R. and Naash, M.I., Properties of the mouse cone-mediated electroretinogram during light adaptation, 9
Pearson, A.D.J., see Lovat, P.E., 109
Perney, P., see Mousseau, D.D., 192

Ikeda, K., Akiyama, H., Kondo, H. and Ikeda, K., Anti-tau-positive glial fibrillary tangles in the brain of postencephalitic parkinsonism of Economo type, 176
Imagawa, M., see Tanaka, H., 63
Inoki, R., see Uchida, Y., 114
Ishii, H., see Yoshida, M., 5

Jaffe, E.H., De Frias, V. and Ibarra, C., Changes in basal and stimulated release of endogenous serotonin from different nuclei of rats subjected to two models of depression, 157
Jellinger, K.A., see Bancher, C., 179
Johnson, J.A., see Tizzano, J.P., 12

Kamino, K., see Tanaka, H., 63
Katao, Y., see Minami, M., 161
Katsumata, S., see Minami, M., 161
Kawahara, H., Yoshida, M., Yokoo, H., Nishi, M. and Tanaka, M., Psychological stress increases serotonin release in the rat amygdala and prefrontal cortex assessed by in vivo microdialysis, 81
Keller, J.T., see Stanley, M.I., 89
Kent, T.A., see Stroemer, R.P., 51
Kiessling, M., see Gass, P., 39
Kim, H.J., see Chung, K., 85
Kimura, J., see Shimohama, S., 183
Kobayashi, H., see Tanaka, H., 63
Kobayashi, J.-i., Ohta, M. and Terada, Y., C fiber generates a slow Na^+ spike in the frog sciatic nerve, 93
Koda, H., see Mizumura, K., 75
Koga, C., see Yoshida, M., 5
Köhler, R., see Campos-Barros, A., 145
Kojima, T., see Tanaka, H., 63
Kondo, H., see Ikeda, K., 176
Krnjević, K., see Godfraind, J.M., 101
Kumazawa, T., see Mizumura, K., 75

Layrargues, G.P., see Mousseau, D.D., 192
Little, J., see Molchan, S.E., 187
Lovat, P.E., Pearson, A.D.J., Malcolm, A. and Redfern, C.P.F., Retinoic acid receptor expression during the in vitro differentiation of human neuroblastoma, 109
Lustig, H.S., see Ahern, K.v.B., 169

Maeda, S., see Uchida, Y., 114
Maekawa, K., see Minami, M., 161
Maeno, H., see Fujita, M., 59
Malcolm, A., see Lovat, P.E., 109
Manji, H., see Molchan, S.E., 187
Martínez, A., see de Rada, O.D., 121
Matsumoto, I., see Pujic, Z., 67
Matsumoto, S., see Shimohama, S., 183
Matsuya, T., see Uchida, Y., 114
May, T., 1-Methyl-4-phenylpyridinium (MPP^+) binds with high affinity to a β-carboline binding site located on monoamine oxidase type A in rat brain, 55
Mechoulam, R., see Nadler, V., 43
Meier, R.K., see Dieringer, N., 129
Meinhold, H., see Campos-Barros, A., 145
Miki, T., see Tanaka, H., 63
Minami, M., Hosoi, Y., Toya, T., Katao, Y., Maekawa, K., Katsumata, S., Yabuuchi, K., Onogi, T. and Satoh, M., In situ hybridization study of κ-opioid receptor mRNA in the rat brain, 161
Miyatake, T., see Tanaka, H., 63
Miyoshi, K., see Shimohama, S., 183
Mizoguchi, K., see Yoshida, M., 5
Mizumura, K., Koda, H. and Kumazawa, T., Augmenting effects of cyclic AMP on the heat response of canine testicular polymodal receptors, 75
Molchan, S.E., Manji, H., Chen, G., Dou, L., Little, J., Potter, W.Z. and Sunderland, T., Effects of chronic lithium treatment on platelet PKC isozymes in Alzheimer's and elderly control subjects, 187
Montuenga, L.M., see de Rada, O.D., 121
Morehead, M., see Sen, S., 117
Mousseau, D.D., Perney, P., Layrargues, G.P. and Butterworth, R.F., Selective loss of pallidal dopamine D_2 receptor density in hepatic encephalopathy, 192
Müller, F., see Campos-Barros, A., 145
Murphy, S., see Sen, S., 117
Murphy, S., see Bruner, G., 105

Na, H.S., see Chung, K., 85
Naash, M.I., see Peachey, N.S., 9
Nachemson, A.K. and Bennett, G.J., Does pain damage spinal cord neurons? Transsynaptic degeneration in rat following a surgical incision, 78
Nadler, V., Mechoulam, R. and Sokolovsky, M., The non-psychotropic cannabinoid (+)-(3*S*,4*S*)-7-hydroxy-Δ^6-tetrahydrocannabinol 1,1-dimethylheptyl (HU-211) attenuates *N*-methyl-D-aspartate receptor-mediated neurotoxicity in primary cultures of rat forebrain, 43
Nakano, I., see Tanaka, H., 63
Naruse, S., see Tanaka, H., 63
Nielander, H.B., French, P., Oestreicher, A.B., Gispen, W.H. and Schotman, P., Spontaneous morphological changes by overexpression of the growth-associated protein B-50/GAP-43 in a PC12 cell line, 46
Nishi, M., see Kawahara, H., 81
Nishimura, M., see Shimohama, S., 183
Nishimura, T., see Fujita, M., 59
Nishino, H., see Czurkó, A., 71
Nowak, J.Z. and Wawrocka, M., Intravitreal application of membrane-permeable analogs of cyclic GMP increases serotonin *N*-acetyltransferase (NAT) activity in retinas of light-exposed chicks: comparison with the effect of cyclic AMP analogs and darkness, 21
Nukina, N., see Tanaka, H., 63

Oestreicher, A.B., see Nielander, H.B., 46
Ohnishi, T., see Uchida, Y., 114
Ohta, M., see Kobayashi, J.-i., 93
Okamoto, K., Hirai, S., Amari, M., Watanabe, M. and Sakurai, A., Bunina bodies in amyotrophic lateral sclerosis immunostained with rabbit anti-cystatin C serum, 125
Onodera, O., see Tanaka, H., 63
Onogi, T., see Minami, M., 161

Pang, C.S., Brown, G.M., Tang, P.L. and Pang, S.F., G-protein linked melatonin binding sites in the chicken lung, 17
Pang, S.F., see Pang, C.S., 17
Panzica, F., see Franceschetti, S., 25
Park, M.J., see Chung, K., 85
Peachey, N.S., Goto, Y., Al-Ubaidi, M.R. and Naash, M.I., Properties of the mouse cone-mediated electroretinogram during light adaptation, 9
Pearson, A.D.J., see Lovat, P.E., 109
Perney, P., see Mousseau, D.D., 192

Perry, G., see Shimohama, S., 183
Phillis, J.W., see Sen, S., 117
Piovacari, G., see Goso, C., 197
Polak, J.M., see de Rada, O.D., 121
Potter, W.Z., see Molchan, S.E., 187
Pujic, Z., Matsumoto, I. and Wilce, P.A., Expression of the gene coding for the NR1 subunit of the NMDA receptor during rat brain development, 67

Rajdev, S. and Reynolds, I.J., Calcium green-5N, a novel fluorescent probe for monitoring high intracellular free Ca^{2+} concentrations associated with glutamate excitotoxicity in cultured rat brain neurons, 149
Redfern, C.P.F., see Lovat, P.E., 109
Reynolds, I.J., see Rajdev, S., 149
Richey, P., see Shimohama, S., 183
Rivero, F.L., see Castellano, M.A., 1
Rodriguez, M., see Castellano, M.A., 1
Rosales, M.G., see Flores, G., 153

Saito, K., see Uchida, Y., 114
Sakaki, Y., see Tanaka, H., 63
Sakurai, A., see Okamoto, K., 125
Sancini, G., see Franceschetti, S., 25
Satoh, M., see Minami, M., 161
Schmidt, R.E., see Dorsey, D.A., 29
Schoepp, D.D., see Tizzano, J.P., 12
Schotman, P., see Nielander, H.B., 46
Seki, K., see Tanaka, H., 63
Sen, S., Goldman, H., Morehead, M., Murphy, S. and Phillis, J.W., Oxypurinol inhibits free radical release from the cerebral cortex of closed head injured rats, 117
Shibata, A., see Tanaka, H., 63
Shimada, S., see Fujita, M., 59
Shimizu, T., see Tanaka, H., 63
Shimohama, S., Perry, G., Richey, P., Takenawa, T., Whitehouse, P.J., Miyoshi, K., Suenaga, T., Matsumoto, S., Nishimura, M. and Kimura, J., Abnormal accumulation of phospholipase C-δ in filamentous inclusions of human neurodegenerative diseases, 183
Sierra, A., see Flores, G., 153
Sokolovsky, M., see Nadler, V., 43
Springall, D.R., see de Rada, O.D., 121
Stanley, M.I., Berger, R.J., Zuccarello, M. and Keller, J.T., Serotonin (5-HT) fibers of the rat dura mater: 5-HT-positive, but not authentic serotoninergic, tryptophan hydroxylase-like fibers, 89
Stroemer, R.P., Kent, T.A. and Hulsebosch, C.E., Acute increase in expression of growth associated protein GAP-43 following cortical ischemia in rat, 51
Suenaga, T., see Shimohama, S., 183
Sunderland, T., see Molchan, S.E., 187
Szallasi, A., see Goso, C., 197

Takenawa, T., see Shimohama, S., 183
Tanaka, H., Naruse, S., Seki, K., Onodera, O., Kobayashi, H., Miyatake, T., Shibata, A., Sakaki, Y., Kamino, K., Miki, T., Nukina, N., Imagawa, M., Nakano, I., Shimizu, T., Kojima, T., Hardy, J. and Tsuji, S., Absence of linkage disequilibrium at amyloid precursor protein gene locus in Japanese familial Alzheimer's disease with ^{717}Val→Ile mutation, 63
Tanaka, M., see Kawahara, H., 81
Tanaka, M., see Yoshida, M., 5
Tanaka, T., see Yoshida, M., 5
Tang, P.L., see Pang, C.S., 17
Terada, Y., see Kobayashi, J.-i., 93
Tizzano, J.P., Griffey, K.I., Johnson, J.A., Fix, A.S., Helton, D.R. and Schoepp, D.D., Intracerebral 1S,3R-1-aminocyclopentane-1,3-dicarboxylic acid (1S,3R-ACPD) produces limbic seizures that are not blocked by ionotropic glutamate receptor antagonists, 12
Tohyama, M., see Fujita, M., 59
Toya, T., see Minami, M., 161
Tsuji, S., see Tanaka, H., 63

Uchida, Y., Saito, K., Ohnishi, T., Maeda, S., Matsuya, T. and Inoki, R., Antagonism by glibenclamide of the effect of morphine in hippocampal preparations, 114
Unsicker, H.-W.K.K., see Bieger, S., 173

Valencia, J., see Flores, G., 153
Valldeoriola, F., see Fueyo, J., 141
Villaro, A.C., see de Rada, O.D., 121

Watanabe, M., see Okamoto, K., 125
Wawrocka, M., see Nowak, J.Z., 21
Wesemann, W., see Campos-Barros, A., 145
Whitehouse, P.J., see Shimohama, S., 183
Wilce, P.A., see Pujic, Z., 67

Yabuuchi, K., see Minami, M., 161
Yokoo, H., see Kawahara, H., 81
Yokoo, H., see Yoshida, M., 5
Yoshida, M., Mizoguchi, K., Yokoo, H., Tanaka, T., Ishii, H., Emoto, H., Koga, C. and Tanaka, M., Muscimol-induced increase in dopamine release and metabolism is not observed in kainic acid-lesioned striatum of conscious rats: an in vivo microdialysis study, 5
Yoshida, M., see Kawahara, H., 81

Zuccarello, M., see Stanley, M.I., 89